

# VECTOR ALGEBRA

## 10.1 Overview

**10.1.1** A quantity that has magnitude as well as direction is called a vector.

**10.1.2** The unit vector in the direction of  $\vec{a}$  is given by  $\frac{\vec{a}}{|\vec{a}|}$  and is represented by  $\hat{a}$ .

**10.1.3** Position vector of a point P ( $x, y, z$ ) is given as  $\vec{OP} = x\hat{i} + y\hat{j} + z\hat{k}$  and its magnitude as  $|\vec{OP}| = \sqrt{x^2 + y^2 + z^2}$ , where O is the origin.

**10.1.4** The scalar components of a vector are its direction ratios, and represent its projections along the respective axes.

**10.1.5** The magnitude  $r$ , direction ratios ( $a, b, c$ ) and direction cosines ( $l, m, n$ ) of any vector are related as:

$$l = \frac{a}{r}, m = \frac{b}{r}, n = \frac{c}{r}.$$

**10.1.6** The sum of the vectors representing the three sides of a triangle taken in order is  $\vec{0}$

**10.1.7** The triangle law of vector addition states that “If two vectors are represented by two sides of a triangle taken in order, then their sum or resultant is given by the third side taken in opposite order”.

### 10.1.8 Scalar multiplication

If  $\vec{a}$  is a given vector and  $\lambda$  a scalar, then  $\lambda\vec{a}$  is a vector whose magnitude is  $|\lambda\vec{a}| = |\lambda||\vec{a}|$ . The direction of  $\lambda\vec{a}$  is same as that of  $\vec{a}$  if  $\lambda$  is positive and, opposite to that of  $\vec{a}$  if  $\lambda$  is negative.

**10.1.9 Vector joining two points**

If  $P_1 (x_1, y_1, z_1)$  and  $P_2 (x_2, y_2, z_2)$  are any two points, then

$$\overrightarrow{P_1P_2} = (x_2 - x_1) \hat{i} + (y_2 - y_1) \hat{j} + (z_2 - z_1) \hat{k}$$

$$|\overrightarrow{P_1P_2}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

**10.1.10 Section formula**

The position vector of a point R dividing the line segment joining the points P and Q whose position vectors are  $\vec{a}$  and  $\vec{b}$

(i) in the ratio  $m : n$  internally, is given by  $\frac{n\vec{a} + m\vec{b}}{m + n}$

(ii) in the ratio  $m : n$  externally, is given by  $\frac{m\vec{b} - n\vec{a}}{m - n}$

**10.1.11** Projection of  $\vec{a}$  along  $\vec{b}$  is  $\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$  and the Projection vector of  $\vec{a}$  along  $\vec{b}$

$$\text{is } \left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \right) \vec{b}.$$

**10.1.12 Scalar or dot product**

The scalar or dot product of two given vectors  $\vec{a}$  and  $\vec{b}$  having an angle  $\theta$  between them is defined as

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$$

**10.1.13 Vector or cross product**

The cross product of two vectors  $\vec{a}$  and  $\vec{b}$  having angle  $\theta$  between them is given as

$$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{n},$$

where  $\hat{n}$  is a unit vector perpendicular to the plane containing  $\vec{a}$  and  $\vec{b}$  and  $\vec{a}, \vec{b}, \hat{n}$  form a right handed system.

**10.1.14** If  $\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$  and  $\vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$  are two vectors and  $\lambda$  is any scalar, then

$$\vec{a} + \vec{b} = (a_1 + b_1)\hat{i} + (a_2 + b_2)\hat{j} + (a_3 + b_3)\hat{k}$$

$$\lambda \vec{a} = (\lambda a_1)\hat{i} + (\lambda a_2)\hat{j} + (\lambda a_3)\hat{k}$$

$$\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = (b_1 c_2 - b_2 c_1)\hat{i} + (a_2 c_1 - c_1 c_2)\hat{j} + (a_1 b_2 - a_2 b_1)\hat{k}$$

Angle between two vectors  $\vec{a}$  and  $\vec{b}$  is given by

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{a_1 b_1 + a_2 b_2 + a_3 b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}}$$

## 10.2 Solved Examples

### Short Answer (S.A.)

**Example 1** Find the unit vector in the direction of the sum of the vectors

$$\vec{a} = 2\hat{i} - \hat{j} + 2\hat{k} \text{ and } \vec{b} = -\hat{i} + \hat{j} + 3\hat{k}.$$

**Solution** Let  $\vec{c}$  denote the sum of  $\vec{a}$  and  $\vec{b}$ . We have

$$\vec{c} = (2\hat{i} - \hat{j} + 2\hat{k}) + (-\hat{i} + \hat{j} + 3\hat{k}) = \hat{i} + 5\hat{k}$$

$$\text{Now } |\vec{c}| = \sqrt{1^2 + 5^2} = \sqrt{26}.$$

Thus, the required unit vector is  $\hat{c} = \frac{\vec{c}}{|\vec{c}|} = \frac{1}{\sqrt{26}}(\hat{i} + 5\hat{k}) = \frac{1}{\sqrt{26}}\hat{i} + \frac{5}{\sqrt{26}}\hat{k}$ .

**Example 2** Find a vector of magnitude 11 in the direction opposite to that of  $\overline{PQ}$ , where P and Q are the points (1, 3, 2) and (-1, 0, 8), respectively.

**Solution** The vector with initial point P(1, 3, 2) and terminal point Q(-1, 0, 8) is given by

$$\overline{PQ} = (-1 - 1)\hat{i} + (0 - 3)\hat{j} + (8 - 2)\hat{k} = -2\hat{i} - 3\hat{j} + 6\hat{k}$$

Thus  $\overline{QP} = -\overline{PQ} = 2\hat{i} + 3\hat{j} - 6\hat{k}$

$$\Rightarrow |\overline{QP}| = \sqrt{2^2 + 3^2 + (-6)^2} = \sqrt{4 + 9 + 36} = \sqrt{49} = 7$$

Therefore, unit vector in the direction of  $\overline{QP}$  is given by

$$\widehat{QP} = \frac{\overline{QP}}{|\overline{QP}|} = \frac{2\hat{i} + 3\hat{j} - 6\hat{k}}{7}$$

Hence, the required vector of magnitude 11 in direction of  $\overline{QP}$  is

$$11 \widehat{QP} = 11 \left( \frac{2\hat{i} + 3\hat{j} - 6\hat{k}}{7} \right) = \frac{22}{7}\hat{i} + \frac{33}{7}\hat{j} - \frac{66}{7}\hat{k}.$$

**Example 3** Find the position vector of a point R which divides the line joining the two points P and Q with position vectors  $\overline{OP} = 2\vec{a} + \vec{b}$  and  $\overline{OQ} = \vec{a} - 2\vec{b}$ , respectively, in the ratio 1:2, (i) internally and (ii) externally.

**Solution** (i) The position vector of the point R dividing the join of P and Q internally in the ratio 1:2 is given by

$$\overline{OR} = \frac{2(2\vec{a} + \vec{b}) + 1(\vec{a} - 2\vec{b})}{1 + 2} = \frac{5\vec{a}}{3}.$$

- (ii) The position vector of the point  $R'$  dividing the join of  $P$  and  $Q$  in the ratio  $1 : 2$  externally is given by

$$\overline{OR'} = \frac{2(2\bar{a} + \bar{b}) - 1(\bar{a} - 2\bar{b})}{2 - 1} = 3\bar{a} + 4\bar{b}.$$

**Example 4** If the points  $(-1, -1, 2)$ ,  $(2, m, 5)$  and  $(3, 11, 6)$  are collinear, find the value of  $m$ .

**Solution** Let the given points be  $A(-1, -1, 2)$ ,  $B(2, m, 5)$  and  $C(3, 11, 6)$ . Then

$$\overline{AB} = (2+1)\hat{i} + (m+1)\hat{j} + (5-2)\hat{k} = 3\hat{i} + (m+1)\hat{j} + 3\hat{k}$$

and  $\overline{AC} = (3+1)\hat{i} + (11+1)\hat{j} + (6-2)\hat{k} = 4\hat{i} + 12\hat{j} + 4\hat{k}.$

Since  $A, B, C$ , are collinear, we have  $\overline{AB} = \lambda \overline{AC}$ , i.e.,

$$(3\hat{i} + (m+1)\hat{j} + 3\hat{k}) = (4\hat{i} + 12\hat{j} + 4\hat{k})$$

$$\Rightarrow 3 = 4\lambda \text{ and } m+1 = 12\lambda$$

Therefore  $m = 8$ .

**Example 5** Find a vector  $\vec{r}$  of magnitude  $3\sqrt{2}$  units which makes an angle of  $\frac{\pi}{4}$  and

$\frac{\pi}{2}$  with  $y$  and  $z$ -axes, respectively.

**Solution** Here  $m = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$  and  $n = \cos \frac{\pi}{2} = 0$ .

Therefore,  $l^2 + m^2 + n^2 = 1$  gives

$$l^2 + \frac{1}{2} + 0 = 1$$

$$\Rightarrow l = \pm \frac{1}{\sqrt{2}}$$

Hence, the required vector  $\vec{r} = 3\sqrt{2} (l\hat{i} + m\hat{j} + n\hat{k})$  is given by

$$\vec{r} = 3\sqrt{2} \left( \pm \frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{j} + 0\hat{k} \right) = \vec{r} = \pm 3\hat{i} + 3\hat{j}.$$

**Example 6** If  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} + \hat{j} - 2\hat{k}$  and  $\vec{c} = \hat{i} + 3\hat{j} - \hat{k}$ , find  $\lambda$  such that  $\vec{a}$  is perpendicular to  $\lambda\vec{b} + \vec{c}$ .

**Solution** We have

$$\begin{aligned} \lambda\vec{b} + \vec{c} &= \lambda(\hat{i} + \hat{j} - 2\hat{k}) + (\hat{i} + 3\hat{j} - \hat{k}) \\ &= (\lambda + 1)\hat{i} + (\lambda + 3)\hat{j} - (2\lambda + 1)\hat{k} \end{aligned}$$

Since  $\vec{a} \perp (\lambda\vec{b} + \vec{c})$ ,  $\vec{a} \cdot (\lambda\vec{b} + \vec{c}) = 0$

$$\begin{aligned} \Rightarrow (2\hat{i} - \hat{j} + \hat{k}) \cdot [(\lambda + 1)\hat{i} + (\lambda + 3)\hat{j} - (2\lambda + 1)\hat{k}] &= 0 \\ \Rightarrow 2(\lambda + 1) - (\lambda + 3) - (2\lambda + 1) &= 0 \\ \Rightarrow \lambda &= -2. \end{aligned}$$

**Example 7** Find all vectors of magnitude  $10\sqrt{3}$  that are perpendicular to the plane of  $\hat{i} + 2\hat{j} + \hat{k}$  and  $-\hat{i} + 3\hat{j} + 4\hat{k}$ .

**Solution** Let  $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$  and  $\vec{b} = -\hat{i} + 3\hat{j} + 4\hat{k}$ . Then

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 1 \\ -1 & 3 & 4 \end{vmatrix} = \hat{i}(8-3) - \hat{j}(4+1) + \hat{k}(3+2) = 5\hat{i} - 5\hat{j} + 5\hat{k}$$

$$\Rightarrow |\vec{a} \times \vec{b}| = \sqrt{(5)^2 + (-5)^2 + (5)^2} = \sqrt{3(5)^2} = 5\sqrt{3}.$$

Therefore, unit vector perpendicular to the plane of  $\vec{a}$  and  $\vec{b}$  is given by

$$\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} = \frac{5\hat{i} - 5\hat{j} + 5\hat{k}}{5\sqrt{3}}$$

Hence, vectors of magnitude of  $10\sqrt{3}$  that are perpendicular to plane of  $\vec{a}$  and  $\vec{b}$

are  $\pm 10\sqrt{3} \left( \frac{5\hat{i} - 5\hat{j} + 5\hat{k}}{5\sqrt{3}} \right)$ , i.e.,  $\pm 10(\hat{i} - \hat{j} + \hat{k})$ .

**Long Answer (L.A.)**

**Example 8** Using vectors, prove that  $\cos(A - B) = \cos A \cos B + \sin A \sin B$ .

**Solution** Let  $\widehat{OP}$  and  $\widehat{OQ}$  be unit vectors making angles  $A$  and  $B$ , respectively, with positive direction of  $x$ -axis. Then  $\angle QOP = A - B$  [Fig. 10.1]

We know  $\widehat{OP} = \overline{OM} + \overline{MP} = \hat{i} \cos A + \hat{j} \sin A$  and  $\widehat{OQ} = \overline{ON} + \overline{NQ} = \hat{i} \cos B + \hat{j} \sin B$ .

By definition  $\widehat{OP} \cdot \widehat{OQ} = |\widehat{OP}| |\widehat{OQ}| \cos(A - B)$

$$= \cos(A - B) \quad \dots (1) \quad (\because |\widehat{OP}| = 1 = |\widehat{OQ}|)$$

In terms of components, we have

$$\begin{aligned} \widehat{OP} \cdot \widehat{OQ} &= (\hat{i} \cos A + \hat{j} \sin A) \cdot (\hat{i} \cos B + \hat{j} \sin B) \\ &= \cos A \cos B + \sin A \sin B \quad \dots (2) \end{aligned}$$

From (1) and (2), we get

$$\cos(A - B) = \cos A \cos B + \sin A \sin B.$$




Fig. 10.1

**Example 9** Prove that in a  $\Delta ABC$ ,  $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$ , where  $a, b, c$  represent the magnitudes of the sides opposite to vertices A, B, C, respectively.

**Solution** Let the three sides of the triangle BC, CA and AB be represented by  $\vec{a}, \vec{b}$  and  $\vec{c}$ , respectively [Fig. 10.2].

We have  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ . i.e.,  $\vec{a} + \vec{b} = -\vec{c}$

which pre cross multiplying by  $\vec{a}$ , and

post cross multiplying by  $\vec{b}$ , gives

$$\vec{a} \times \vec{b} = \vec{c} \times \vec{a}$$

and  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c}$

respectively. Therefore,

$$\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$$

$$\Rightarrow |\vec{a} \times \vec{b}| = |\vec{b} \times \vec{c}| = |\vec{c} \times \vec{a}|$$

$$\Rightarrow |\vec{a}| |\vec{b}| \sin(\pi - C) = |\vec{b}| |\vec{c}| \sin(\pi - A) = |\vec{c}| |\vec{a}| \sin(\pi - B)$$

$$\Rightarrow ab \sin C = bc \sin A = ca \sin B$$

Dividing by  $abc$ , we get

$$\frac{\sin C}{c} = \frac{\sin A}{a} = \frac{\sin B}{b} \text{ i.e. } \frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

### Objective Type Questions

Choose the correct answer from the given four options in each of the Examples 10 to 21.

**Example 10** The magnitude of the vector  $6\hat{i} + 2\hat{j} + 3\hat{k}$  is




Fig. 10.2



- (A) 5                      (B) 7                      (C) 12                      (D) 1

**Solution** (B) is the correct answer.

**Example 11** The position vector of the point which divides the join of points with position vectors  $\vec{a} + \vec{b}$  and  $2\vec{a} - \vec{b}$  in the ratio 1 : 2 is

- (A)  $\frac{3\vec{a} + 2\vec{b}}{3}$                       (B)  $\vec{a}$                       (C)  $\frac{5\vec{a} - \vec{b}}{3}$                       (D)  $\frac{4\vec{a} + \vec{b}}{3}$

**Solution** (D) is the correct answer. Applying section formula the position vector of the required point is

$$\frac{2(\vec{a} + \vec{b}) + 1(2\vec{a} - \vec{b})}{2 + 1} = \frac{4\vec{a} + \vec{b}}{3}$$

**Example 12** The vector with initial point P (2, -3, 5) and terminal point Q(3, -4, 7) is

- (A)  $\hat{i} - \hat{j} + 2\hat{k}$                       (B)  $5\hat{i} - 7\hat{j} + 12\hat{k}$   
 (C)  $-\hat{i} + \hat{j} - 2\hat{k}$                       (D) None of these

**Solution** (A) is the correct answer.

**Example 13** The angle between the vectors  $\hat{i} - \hat{j}$  and  $\hat{j} - \hat{k}$  is

- (A)  $\frac{\pi}{3}$                       (B)  $\frac{2\pi}{3}$                       (C)  $\frac{-\pi}{3}$                       (D)  $\frac{5\pi}{6}$

**Solution** (B) is the correct answer. Apply the formula  $\cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| \cdot |\vec{b}|}$ .

**Example 14** The value of  $\lambda$  for which the two vectors  $2\hat{i} - \hat{j} + 2\hat{k}$  and  $3\hat{i} + \lambda\hat{j} + \hat{k}$  are perpendicular is

- (A) 2                      (B) 4                      (C) 6                      (D) 8

**Solution** (D) is the correct answer.

**Example 15** The area of the parallelogram whose adjacent sides are  $\hat{i} + \hat{k}$  and  $2\hat{i} + \hat{j} + \hat{k}$  is

- (A)  $\sqrt{2}$  (B)  $\sqrt{3}$  (C) 3 (D) 4

**Solution** (B) is the correct answer. Area of the parallelogram whose adjacent sides are  $\vec{a}$  and  $\vec{b}$  is  $|\vec{a} \times \vec{b}|$ .

**Example 16** If  $|\vec{a}| = 8$ ,  $|\vec{b}| = 3$  and  $|\vec{a} \times \vec{b}| = 12$ , then value of  $\vec{a} \cdot \vec{b}$  is

- (A)  $6\sqrt{3}$  (B)  $8\sqrt{3}$  (C)  $12\sqrt{3}$  (D) None of these

**Solution** (C) is the correct answer. Using the formula  $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| |\sin \theta|$ , we get

$$\theta = \pm \frac{\pi}{6}.$$

Therefore,  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta = 8 \times 3 \times \frac{\sqrt{3}}{2} = 12\sqrt{3}$ .

**Example 17** The 2 vectors  $\hat{j} + \hat{k}$  and  $3\hat{i} - \hat{j} + 4\hat{k}$  represents the two sides AB and AC, respectively of a  $\Delta ABC$ . The length of the median through A is

- (A)  $\frac{\sqrt{34}}{2}$  (B)  $\frac{\sqrt{48}}{2}$  (C)  $\sqrt{18}$  (D) None of these

**Solution** (A) is the correct answer. Median  $\overline{AD}$  is given by

$$|\overline{AD}| = \frac{1}{2} |3\hat{i} + \hat{j} + 5\hat{k}| = \frac{\sqrt{34}}{2}$$

**Example 18** The projection of vector  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$  along  $\vec{b} = \hat{i} + 2\hat{j} + 2\hat{k}$  is

- (A)  $\frac{2}{3}$                       (B)  $\frac{1}{3}$                       (C) 2                      (D)  $\sqrt{6}$

**Solution** (A) is the correct answer. Projection of a vector  $\vec{a}$  on  $\vec{b}$  is

$$\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{(2\hat{i} - \hat{j} + \hat{k}) \cdot (\hat{i} + 2\hat{j} + 2\hat{k})}{\sqrt{1+4+4}} = \frac{2}{3}.$$

**Example 19** If  $\vec{a}$  and  $\vec{b}$  are unit vectors, then what is the angle between  $\vec{a}$  and  $\vec{b}$  for  $\sqrt{3}\vec{a} - \vec{b}$  to be a unit vector?

- (A)  $30^\circ$                       (B)  $45^\circ$                       (C)  $60^\circ$                       (D)  $90^\circ$

**Solution** (A) is the correct answer. We have

$$(\sqrt{3}\vec{a} - \vec{b})^2 = 3\vec{a}^2 + \vec{b}^2 - 2\sqrt{3}\vec{a} \cdot \vec{b}$$

$$\Rightarrow \vec{a} \cdot \vec{b} = \frac{\sqrt{3}}{2} \Rightarrow \cos\theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = 30^\circ.$$

**Example 20** The unit vector perpendicular to the vectors  $\hat{i} - \hat{j}$  and  $\hat{i} + \hat{j}$  forming a right handed system is

- (A)  $\hat{k}$                       (B)  $-\hat{k}$                       (C)  $\frac{\hat{i} - \hat{j}}{\sqrt{2}}$                       (D)  $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$

**Solution** (A) is the correct answer. Required unit vector is  $\frac{(\hat{i} - \hat{j}) \times (\hat{i} + \hat{j})}{|(\hat{i} - \hat{j}) \times (\hat{i} + \hat{j})|} = \frac{2\hat{k}}{2} = \hat{k}$ .

**Example 21** If  $|\vec{a}|=3$  and  $-1 \leq k \leq 2$ , then  $|k\vec{a}|$  lies in the interval

- (A)  $[0, 6]$                       (B)  $[-3, 6]$                       (C)  $[3, 6]$                       (D)  $[1, 2]$

**Solution** (A) is the correct answer. The smallest value of  $|k\vec{a}|$  will exist at numerically smallest value of  $k$ , i.e., at  $k = 0$ , which gives  $|k\vec{a}| = |k||\vec{a}| = 0 \times 3 = 0$

The numerically greatest value of  $k$  is 2 at which  $|k\vec{a}| = 6$ .

### 10.3 EXERCISE

#### Short Answer (S.A.)

- Find the unit vector in the direction of sum of vectors  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$  and  $\vec{b} = 2\hat{j} + \hat{k}$ .
- If  $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$  and  $\vec{b} = 2\hat{i} + \hat{j} - 2\hat{k}$ , find the unit vector in the direction of
  - $6\vec{b}$
  - $2\vec{a} - \vec{b}$
- Find a unit vector in the direction of  $\overline{PQ}$ , where P and Q have co-ordinates (5, 0, 8) and (3, 3, 2), respectively.
- If  $\vec{a}$  and  $\vec{b}$  are the position vectors of A and B, respectively, find the position vector of a point C in BA produced such that  $BC = 1.5 BA$ .
- Using vectors, find the value of  $k$  such that the points  $(k, -10, 3)$ ,  $(1, -1, 3)$  and  $(3, 5, 3)$  are collinear.
- A vector  $\vec{r}$  is inclined at equal angles to the three axes. If the magnitude of  $\vec{r}$  is  $2\sqrt{3}$  units, find  $\vec{r}$ .
- A vector  $\vec{r}$  has magnitude 14 and direction ratios 2, 3, -6. Find the direction cosines and components of  $\vec{r}$ , given that  $\vec{r}$  makes an acute angle with  $x$ -axis.
- Find a vector of magnitude 6, which is perpendicular to both the vectors  $2\hat{i} - \hat{j} + 2\hat{k}$  and  $4\hat{i} - \hat{j} + 3\hat{k}$ .
- Find the angle between the vectors  $2\hat{i} - \hat{j} + \hat{k}$  and  $3\hat{i} + 4\hat{j} - \hat{k}$ .
- If  $\vec{a} + \vec{b} + \vec{c} = 0$ , show that  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$ . Interpret the result geometrically?
- Find the sine of the angle between the vectors  $\vec{a} = 3\hat{i} + \hat{j} + 2\hat{k}$  and  $\vec{b} = 2\hat{i} - 2\hat{j} + 4\hat{k}$ .

12. If A, B, C, D are the points with position vectors  $\hat{i} + \hat{j} - \hat{k}$ ,  $2\hat{i} - \hat{j} + 3\hat{k}$ ,  $2\hat{i} - 3\hat{k}$ ,  $3\hat{i} - 2\hat{j} + \hat{k}$ , respectively, find the projection of  $\overline{AB}$  along  $\overline{CD}$ .
13. Using vectors, find the area of the triangle ABC with vertices A(1, 2, 3), B(2, -1, 4) and C(4, 5, -1).
14. Using vectors, prove that the parallelogram on the same base and between the same parallels are equal in area.

**Long Answer (L.A.)**

15. Prove that in any triangle ABC,  $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$ , where  $a, b, c$  are the magnitudes of the sides opposite to the vertices A, B, C, respectively.
16. If  $\vec{a}, \vec{b}, \vec{c}$  determine the vertices of a triangle, show that  $\frac{1}{2}[\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + \vec{a} \times \vec{b}]$  gives the vector area of the triangle. Hence deduce the condition that the three points  $\vec{a}, \vec{b}, \vec{c}$  are collinear. Also find the unit vector normal to the plane of the triangle.
17. Show that area of the parallelogram whose diagonals are given by  $\vec{a}$  and  $\vec{b}$  is  $\frac{|\vec{a} \times \vec{b}|}{2}$ . Also find the area of the parallelogram whose diagonals are  $2\hat{i} - \hat{j} + \hat{k}$  and  $\hat{i} + 3\hat{j} - \hat{k}$ .
18. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  and  $\vec{b} = \hat{j} - \hat{k}$ , find a vector  $\vec{c}$  such that  $\vec{a} \times \vec{c} = \vec{b}$  and  $\vec{a} \cdot \vec{c} = 3$ .

**Objective Type Questions**

Choose the correct answer from the given four options in each of the Exercises from 19 to 33 (M.C.Q)

19. The vector in the direction of the vector  $\hat{i} - 2\hat{j} + 2\hat{k}$  that has magnitude 9 is
- |     |                                    |     |   |
|-----|------------------------------------|-----|---|
| (A) | $\hat{i} - 2\hat{j} + 2\hat{k}$    | (B) | $\frac{\hat{i} - 2\hat{j} + 2\hat{k}}{3}$ |
| (C) | $3(\hat{i} - 2\hat{j} + 2\hat{k})$ | (D) | $9(\hat{i} - 2\hat{j} + 2\hat{k})$        |

20. The position vector of the point which divides the join of points  $2\vec{a} - 3\vec{b}$  and  $\vec{a} + \vec{b}$  in the ratio 3 : 1 is

(A)  $\frac{3\vec{a} - 2\vec{b}}{2}$  (B)  $\frac{7\vec{a} - 8\vec{b}}{4}$  (C)  $\frac{3\vec{a}}{4}$  (D)  $\frac{5\vec{a}}{4}$

21. The vector having initial and terminal points as (2, 5, 0) and (-3, 7, 4), respectively is

(A)  $-\hat{i} + 12\hat{j} + 4\hat{k}$  (B)  $5\hat{i} + 2\hat{j} - 4\hat{k}$   
 (C)  $-5\hat{i} + 2\hat{j} + 4\hat{k}$  (D)  $\hat{i} + \hat{j} + \hat{k}$

22. The angle between two vectors  $\vec{a}$  and  $\vec{b}$  with magnitudes  $\sqrt{3}$  and 4, respectively, and  $\vec{a} \cdot \vec{b} = 2\sqrt{3}$  is

(A)  $\frac{\pi}{6}$  (B)  $\frac{\pi}{3}$  (C)  $\frac{\pi}{2}$  (D)  $\frac{5\pi}{2}$

23. Find the value of  $\lambda$  such that the vectors  $\vec{a} = 2\hat{i} + \lambda\hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$  are orthogonal

(A) 0 (B) 1 (C)  $\frac{3}{2}$  (D)  $-\frac{5}{2}$

24. The value of  $\lambda$  for which the vectors  $3\hat{i} - 6\hat{j} + \hat{k}$  and  $2\hat{i} - 4\hat{j} + \lambda\hat{k}$  are parallel is

(A)  $\frac{2}{3}$  (B)  $\frac{3}{2}$  (C)  $\frac{5}{2}$  (D)  $\frac{2}{5}$

25. The vectors from origin to the points A and B are  $\vec{a} = 2\hat{i} - 3\hat{j} + 2\hat{k}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} + \hat{k}$ , respectively, then the area of triangle OAB is

(A) 340 (B)  $\sqrt{25}$  (C)  $\sqrt{229}$  (D)  $\frac{1}{2}\sqrt{229}$

26. For any vector  $\vec{a}$ , the value of  $(\vec{a} \times \hat{i})^2 + (\vec{a} \times \hat{j})^2 + (\vec{a} \times \hat{k})^2$  is equal to  
 (A)  $\vec{a}^2$  (B)  $3\vec{a}^2$  (C)  $4\vec{a}^2$  (D)  $2\vec{a}^2$
27. If  $|\vec{a}| = 10$ ,  $|\vec{b}| = 2$  and  $\vec{a} \cdot \vec{b} = 12$ , then value of  $|\vec{a} \times \vec{b}|$  is  
 (A) 5 (B) 10 (C) 14 (D) 16
28. The vectors  $\lambda \hat{i} + \hat{j} + 2\hat{k}$ ,  $\hat{i} + \lambda \hat{j} - \hat{k}$  and  $2\hat{i} - \hat{j} + \lambda \hat{k}$  are coplanar if  
 (A)  $\lambda = -2$  (B)  $\lambda = 0$  (C)  $\lambda = 1$  (D)  $\lambda = -1$
29. If  $\vec{a}, \vec{b}, \vec{c}$  are unit vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , then the value of  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$  is  
 (A) 1 (B) 3 (C)  $-\frac{3}{2}$  (D) None of these
30. Projection vector of  $\vec{a}$  on  $\vec{b}$  is  
 (A)  $\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \right) \vec{b}$  (B)  $\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$  (C)  $\frac{\vec{a} \cdot \vec{b}}{|\vec{a}|}$  (D)  $\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|^2} \right) \hat{b}$
31. If  $\vec{a}, \vec{b}, \vec{c}$  are three vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $|\vec{a}| = 2$ ,  $|\vec{b}| = 3$ ,  $|\vec{c}| = 5$ , then value of  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$  is  
 (A) 0 (B) 1 (C) -19 (D) 38
32. If  $|\vec{a}| = 4$  and  $-3 \leq \lambda \leq 2$ , then the range of  $|\lambda \vec{a}|$  is  
 (A)  $[0, 8]$  (B)  $[-12, 8]$  (C)  $[0, 12]$  (D)  $[8, 12]$
33. The number of vectors of unit length perpendicular to the vectors  $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$  and  $\vec{b} = \hat{j} + \hat{k}$  is  
 (A) one (B) two (C) three (D) infinite

Fill in the blanks in each of the Exercises from 34 to 40.

34. The vector  $\vec{a} + \vec{b}$  bisects the angle between the non-collinear vectors  $\vec{a}$  and  $\vec{b}$  if \_\_\_\_\_

35. If  $\vec{r} \cdot \vec{a} = 0$ ,  $\vec{r} \cdot \vec{b} = 0$ , and  $\vec{r} \cdot \vec{c} = 0$  for some non-zero vector  $\vec{r}$ , then the value of  $\vec{a} \cdot (\vec{b} \times \vec{c})$  is \_\_\_\_\_.
36. The vectors  $\vec{a} = 3\hat{i} - 2\hat{j} + 2\hat{k}$  and  $\vec{b} = -\hat{i} - 2\hat{k}$  are the adjacent sides of a parallelogram. The acute angle between its diagonals is \_\_\_\_\_.
37. The values of  $k$  for which  $|k\vec{a}| < |\vec{a}|$  and  $k\vec{a} + \frac{1}{2}\vec{a}$  is parallel to  $\vec{a}$  holds true are \_\_\_\_\_.
38. The value of the expression  $|\vec{a} \times \vec{b}|^2 + (\vec{a} \cdot \vec{b})^2$  is \_\_\_\_\_.
39. If  $|\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = 144$  and  $|\vec{a}| = 4$ , then  $|\vec{b}|$  is equal to \_\_\_\_\_.
40. If  $\vec{a}$  is any non-zero vector, then  $(\vec{a} \cdot \hat{i})\hat{i} + (\vec{a} \cdot \hat{j})\hat{j} + (\vec{a} \cdot \hat{k})\hat{k}$  equals \_\_\_\_\_.

State **True** or **False** in each of the following Exercises.

41. If  $|\vec{a}| = |\vec{b}|$ , then necessarily it implies  $\vec{a} = \pm \vec{b}$ .
42. Position vector of a point P is a vector whose initial point is origin.
43. If  $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$ , then the vectors  $\vec{a}$  and  $\vec{b}$  are orthogonal.
44. The formula  $(\vec{a} + \vec{b})^2 = \vec{a}^2 + \vec{b}^2 + 2\vec{a} \times \vec{b}$  is valid for non-zero vectors  $\vec{a}$  and  $\vec{b}$ .
45. If  $\vec{a}$  and  $\vec{b}$  are adjacent sides of a rhombus, then  $\vec{a} \cdot \vec{b} = 0$ .

